## DHANUSH HEALTHCARE

Meru

### **RD Result Report**

### **IMPORTANT INFORMATION:**

- 1. The VIDAS® Total IgE (IGE) is an automated quantitative enzyme linked fluorescent immunoassay (ELFA) for the determination of total IgE concentration in serum on plasma.
- 2. The VIDAS Total IgE assay is intended for use as an aid in the diagnosis of allergic diseases such as such as allergic rhinitis, allergic asthma, and atopic dermatitis Measurement of IgE serum levels can be important in the diagnosis and treatment of these disorders.
- 3. In most nonatopic patients, IgE serum levels are relatively low However, certain parasitic or helminth infections have been associated with elevated IgE levels due to IgE sensitization of macrophages, eosinophils, and other inflammatory cells.
- 4. Elevated IgE levels have also been reported in some forms of glomerular disease and in some immunodeficiency syndromes such as Wiskott- Aldrich syndrome.
- 5. The detection limit for the VIDAS Total IgE assay is 0.5 kIU/l.
- 6. Expected values in Adults: Among a group of healthy people free from any allergic manifestation, 95 % of obtained values are under 120 kIU/l Children.
- 7. Expected values inchildren: These results are given by age(please see below)-

Age group	0 to 20 kIU/l	21 to 50 kIU/l	51 to 10 kIU/l	>100 kIU/l
≤ <b>10</b> months n =93	70.0 %	20.4	6.4 %	3.2 %
2years n =86	31.4 %	23.2 %	18.6 %	<b>26.7</b> %
4years n =49	20.4	20.4	16.3 %	<b>42.9</b> %

# **DHANUSH HEALTHCARE**

Meru

### **RD Result Report**

5 - 10years n =27	11.1	22.2 %	22.2 %	<b>44.5</b> %
>10years n =11	27.3 %	9.0	27.3 %	<b>36.4</b> %

### **LIMITATIONS**

1. Low total IgE concentrations do not necessarily indicate the absence of allergies. Some patients may have a low total IgE but a high concentration of specific IgE antibody.

# **Authorized By**

Dr. Saurabh G Agarwal

**END OF REPORT**